

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LASTACAFT™ safely and effectively. See full prescribing information for LASTACAFT™.

LASTACAFT™ (alcaftadine ophthalmic solution)

Initial U.S. Approval: 2010

INDICATIONS AND USAGE

LASTACAFT™ is a H₁ histamine receptor antagonist indicated for the prevention of itching associated with allergic conjunctivitis (1)

DOSAGE AND ADMINISTRATION

Instill one drop in each eye once daily. (2)

DOSAGE FORMS AND STRENGTHS

Ophthalmic solution containing alcaftadine, 0.25% (2.5 mg/mL) (3)

WARNINGS AND PRECAUTIONS

- To minimize the risk of contamination, do not touch dropper tip to any surface. Keep bottle tightly closed when not in use. (5.1)

- LASTACAFT™ should not be used to treat contact lens-related irritation. (5.2)
- Remove contact lenses prior to instillation of LASTACAFT™. (5.2)

ADVERSE REACTIONS

The most common ocular adverse reactions, occurring in < 4% of LASTACAFT™-treated eyes, were eye irritation, burning and/or stinging on instillation, eye redness, and eye pruritus. (6.1)

The most common non-ocular adverse reactions, occurring in < 3% of subjects with LASTACAFT™-treated eyes, were nasopharyngitis, headache and influenza (6.2)

To report SUSPECTED ADVERSE REACTIONS, contact Vistakon Pharmaceuticals, LLC at 1-800-523-6225 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

To report SUSPECTED ADVERSE REACTIONS, contact at or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

See 17 for PATIENT COUNSELING INFORMATION

Revised: 08/2010

FULL PRESCRIBING INFORMATION: CONTENTS *

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Contamination of Tip and Solution
- 5.2 Contact Lens Use
- 5.3 Topical Ophthalmic Use Only

6 ADVERSE REACTIONS

- 6.1 Ocular Adverse Reactions
- 6.2 Non-ocular Adverse Reactions

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Sterility of Dropper Tip
- 17.2 Concomitant Use of Contact Lenses
- 17.3 Topical Ophthalmic Use only

PRINCIPAL DISPLAY PANEL - 3 ML BOTTLE CARTON

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

LASTACAFT™ is a H₁ histamine receptor antagonist indicated for the prevention of itching associated with allergic conjunctivitis.

2 DOSAGE AND ADMINISTRATION

Instill one drop in each eye once daily.

3 DOSAGE FORMS AND STRENGTHS

Topical ophthalmic solution containing alcaftadine, 0.25% (2.5 mg/mL).

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Contamination of Tip and Solution

To minimize contaminating the dropper tip and solution, care should be taken not to touch the eyelids or surrounding areas with the dropper tip of the bottle. Keep bottle tightly closed when not in use.

5.2 Contact Lens Use

Patients should be advised not to wear a contact lens if their eye is red. LASTACAFT™ should not be used to treat contact lens-related irritation.

LASTACAFT™ should not be instilled while wearing contact lenses. Remove contact lenses prior to instillation of LASTACAFT™. The preservative in LASTACAFT™, benzalkonium chloride, may be absorbed by soft contact lenses. Lenses may be reinserted after 10 minutes following administration of LASTACAFT™.

5.3 Topical Ophthalmic Use Only

LASTACAFT™ is for topical ophthalmic use only.

6 ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

6.1 Ocular Adverse Reactions

The most frequent ocular adverse reactions, occurring in < 4% of LASTACAFT™-treated eyes, were eye irritation, burning and/or stinging upon instillation, eye redness and eye pruritus.

6.2 Non-ocular Adverse Reactions

The most frequent non-ocular adverse reactions, occurring in < 3% of subjects with LASTACAFT™-treated eyes, were nasopharyngitis, headache and influenza. Some of these events were similar to the underlying disease being studied.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B. Reproduction studies performed in rats and rabbits revealed no evidence of impaired female reproduction or harm to the fetus due to alcaftadine. Oral doses in rats and rabbits of 20 and 80 mg/kg/day, respectively, produced plasma exposure levels approximately 200 and 9000 times the plasma exposure at the recommended human ocular dose. There are however, no adequate and well controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

8.3 Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when LASTACAFT™ is administered to a nursing woman.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients below the age of 2 years have not been established.

8.5 Geriatric Use

No overall differences in safety or effectiveness were observed between elderly and younger subjects.

11 DESCRIPTION

LASTACAFT™ is a sterile, topically administered H₁ receptor antagonist containing alcaftadine for ophthalmic use. Alcaftadine is a white to yellow powder with an empirical formula of C₁₉H₂₁N₃O and a molecular weight of 307.39.

Contains:

Active: alcaftadine 0.25% (2.5 mg/mL)

Preservative: benzalkonium chloride 0.005%

Inactives: edetate disodium, monobasic sodium phosphate, purified water, sodium chloride, sodium hydroxide and/or hydrochloric acid (to adjust pH)

Chemical Name: 6,11-dihydro-11-(1-methyl-4-piperidinylidene)-5H-imidazo[2,1-b] [3] benzazepine-3-carboxaldehyde

Structural Formula:



The drug product has a pH of approximately 7 and an osmolality of approximately 290 mOsm/kg.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Alcaftadine is a H₁ histamine receptor antagonist and inhibitor of the release of histamine from mast cells. Decreased chemotaxis and inhibition of eosinophil activation has also been demonstrated.

12.3 Pharmacokinetics

Absorption

Following bilateral topical ocular administration of alcaftadine ophthalmic solution, 0.25%, the mean plasma C_{max} of alcaftadine was approximately 60 pg/mL and the median T_{max} occurred at 15 minutes. Plasma concentrations of alcaftadine were below the lower limit of quantification (10 pg/mL) by 3 hours after dosing. The mean C_{max} of the active carboxylic acid metabolite was approximately 3 ng/mL and occurred at 1 hour after dosing. Plasma concentrations of the carboxylic acid metabolite were below the lower limit of quantification (100 pg/mL) by 12 hours after dosing. There was no indication of systemic accumulation or changes in plasma exposure of alcaftadine or the active metabolite following daily topical ocular administration.

Distribution

The protein binding of alcaftadine and the active metabolite are 39.2% and 62.7%, respectively.

Metabolism

The metabolism of alcaftadine is mediated by non-CYP450 cytosolic enzymes to the active carboxylic acid metabolite.

Excretion

The elimination half-life of the carboxylic acid metabolite is approximately 2 hours following topical ocular administration. Based on data following oral administration of alcaftadine, the carboxylic acid metabolite is primarily eliminated unchanged in the urine.

In vitro studies showed that neither alcaftadine nor the carboxylic acid metabolite substantially inhibited reactions catalyzed by major CYP450 enzymes.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Alcaftadine was not mutagenic or genotoxic in the Ames test, the mouse lymphoma assay or the mouse micronucleus assay. Alcaftadine was found to have no effect on fertility of male and female rats at oral doses up to 20 mg/kg/day (approximately 200 times the plasma exposure at the recommended human ocular dose).

14 CLINICAL STUDIES

Clinical efficacy was evaluated in conjunctival allergen challenge (CAC) studies. LASTACAFT™ was more effective than its vehicle in preventing ocular itching in patients with allergic conjunctivitis induced by an ocular allergen challenge, both at 3 minutes post-dosing and at 16 hours post-dosing of LASTACAFT™.

The safety of LASTACAFT™ was evaluated in a randomized clinical study of 909 subjects over a period of 6 weeks.

16 HOW SUPPLIED/STORAGE AND HANDLING

LASTACAFT™ (alcaftadine ophthalmic solution) 0.25% is supplied in an opaque, white low-density polyethylene bottle with a white polypropylene cap.

3 mL fill in 5 mL bottle (NDC 68669-412-03)

Storage: Store at 15-25°C (59-77°F)

17 PATIENT COUNSELING INFORMATION

17.1 Sterility of Dropper Tip

Patients should be advised to not touch dropper tip to any surface, as this may contaminate the contents.

17.2 Concomitant Use of Contact Lenses

Patients should be advised not to wear a contact lens if their eye is red. Patients should be advised that LASTACAFT™ should not be used to treat contact lens-related irritation. Patients should also be advised to remove contact lenses prior to instillation of LASTACAFT™. The preservative in LASTACAFT™, benzalkonium chloride, may be absorbed by soft contact lenses. Lenses may be reinserted after 10 minutes following administration of LASTACAFT™.

17.3 Topical Ophthalmic Use only

For topical ophthalmic administration only.

Manufactured for Vistakon Pharmaceuticals, LLC

Jacksonville, FL 32256 USA

PRINCIPAL DISPLAY PANEL - 3 ML BOTTLE CARTON

NDC 68669-412-03

LASTACAFT™

(alcaftadine ophthalmic
solution) 0.25%

3 mL

Sterile

VISTAKON® PHARMACEUTICALS, LLC

